

# Advisory Committee on Immunization Practices Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger — United States, 2017

Candice L. Robinson, MD<sup>1</sup>; José R. Romero, MD<sup>2</sup>; Allison Kempe, MD<sup>3</sup>; Cynthia Pellegrini<sup>4</sup>; Advisory Committee on Immunization Practices (ACIP) Child/Adolescent Immunization Work Group

On February 7, 2017, this report was posted as an MMWR Early Release on the MMWR website (<https://www.cdc.gov/mmwr>).

In October 2016, the Advisory Committee on Immunization Practices (ACIP) approved the Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger—United States, 2017. The 2017 child and adolescent immunization schedule summarizes ACIP recommendations, including several changes from the 2016 immunization schedules, in three figures, and footnotes for the figures. These documents can be found on the CDC immunization schedule website (<https://www.cdc.gov/vaccines/schedules/index.html>). These immunization schedules are approved by ACIP (<https://www.cdc.gov/vaccines/acip/index.html>), the American Academy of Pediatrics (<https://www.aap.org>), the American Academy of Family Physicians (<https://www.aafp.org>), and the American College of Obstetricians and Gynecologists (<http://www.acog.org>). Health care providers are advised to use the figures and the combined footnotes together. The full ACIP recommendations for each vaccine, including contraindications and precautions, can be found at <https://www.cdc.gov/vaccines/hcp/acip-recs/index.html>. Providers should be aware that changes in recommendations for specific vaccines can

occur between annual updates to the childhood/adolescent immunization schedules. If errors or omissions are discovered within the child and adolescent schedule, CDC posts revised versions on the CDC immunization schedule website.\*

Printable versions of the 2017 immunization schedules for children and adolescents aged 18 years or younger also are available at the website and ordering instructions for laminated versions and easy-to-read versions for parents also are available at the immunization schedule website.

For further guidance on the use of each vaccine included in the schedules, including contraindications and precautions, health care providers are referred to the respective ACIP vaccine recommendations at <https://www.cdc.gov/vaccines/hcp/acip-recs/index.html>.

## Changes in the 2017 Child and Adolescent Immunization Schedule

Changes in the 2017 immunization schedules for children and adolescents aged 18 years or younger include new or revised ACIP recommendations for influenza (1); human papillomavirus (2); hepatitis B (3); *Haemophilus influenzae* type B (4); pneumococcal; meningococcal (5,6); and diphtheria and tetanus toxoids and acellular pertussis (7) vaccines.

**Figure 1.** Changes to the 2017 figure from the 2016 schedule<sup>†</sup> are as follows:

- The 16-year age column has been separated from the 17–18-year age column to highlight the need for a meningococcal conjugate vaccine booster dose at age 16 years.
- Live attenuated influenza vaccine (LAIV) has been removed from the influenza row.
- A blue bar was added for human papillomavirus vaccine (HPV) for children aged 9–10 years, indicating that persons in this age group may be vaccinated (even in the absence of a high-risk condition).

*Recommendations for routine use of vaccines in children, adolescents and adults are developed by the Advisory Committee on Immunization Practices (ACIP). ACIP is chartered as a federal advisory committee to provide expert external advice and guidance to the Director of the Centers for Disease Control and Prevention (CDC) on use of vaccines and related agents for the control of vaccine-preventable diseases in the civilian population of the United States. Recommendations for routine use of vaccines in children and adolescents are harmonized to the greatest extent possible with recommendations made by the American Academy of Pediatrics (AAP), the American Academy of Family Physicians (AAFP), and the American College of Obstetricians and Gynecologists (ACOG). Recommendations for routine use of vaccines in adults are harmonized with recommendations of AAFP, ACOG, and the American College of Physicians (ACP). ACIP recommendations approved by the CDC Director become agency guidelines on the date published in the Morbidity and Mortality Weekly Report (MMWR). Additional information about ACIP is available at <https://www.cdc.gov/vaccines/acip>.*

\* CDC encourages organizations that previously have relied on copying the schedules to their websites instead to use syndication as a more reliable method for displaying the most current and accurate immunization schedules on an organization's website. Use of content syndication requires a one-time step that ensures an organization's website displays current schedules as soon as they are published or revised; instructions for the syndication code are available on CDC's website (<https://www.cdc.gov/vaccines/schedules/syndicate.html>). CDC also offers technical assistance for implementing this form of content syndication (e-mail request to [ncirdwebteam@cdc.gov](mailto:ncirdwebteam@cdc.gov)).

<sup>†</sup> Past immunization schedules are available at <https://www.cdc.gov/vaccines/schedules/past.html>.

**Figure 3.** A new figure, “Figure 3. Vaccines that might be indicated for children and adolescents aged 18 years or younger based on medical indications,” has been added. The purpose of this figure is to do the following:

- Demonstrate most children with medical conditions can (and should) be vaccinated according to the routine child/adolescent immunization schedule.
- Indicate when a medical condition is a precaution or contraindication to vaccination.
- Indicate when additional doses of vaccines may be necessary because of a child’s or adolescent’s medical condition. Providers should consult the relevant footnotes for additional information.

**Footnotes.** Changes to the footnotes for the figures are as follows:

- The Hepatitis B vaccine (HepB) footnote was revised to reflect that the birth dose of HepB should be administered within 24 hours of birth.
- The diphtheria and tetanus toxoids and acellular pertussis vaccine (DTaP) footnote was revised to more clearly present recommendations following an inadvertently early administered fourth dose of DTaP.
- Within the *Haemophilus influenzae* type b vaccine (Hib) footnote, Comvax was removed from the routine vaccination portion of footnote. This vaccine has been removed from the market, and all available doses have expired. Additionally, Hiberix has been added to the list of vaccines that may be used for the primary vaccination series.
- Within the pneumococcal vaccine footnote, references to 7-valent pneumococcal conjugate vaccine (PCV7) have been removed. All healthy children who might have received PCV7 as part of a primary series have now aged out of the recommendation for pneumococcal vaccine.
- The influenza vaccine footnote has been updated to indicate that LAIV should not be used during the 2016–2017 influenza season.
- The meningococcal vaccines footnote has been updated to include recommendations for meningococcal vaccination of children with human immunodeficiency virus (HIV) infection and to reflect recommendations for the use of a 2-dose Trumenba (meningococcal B vaccine) schedule.
- The tetanus and diphtheria toxoids and acellular pertussis vaccine (Tdap) footnote for vaccination of pregnant adolescents between gestational weeks 27–36 has been updated to reflect a preference for vaccination earlier during this period. Currently available data suggest that vaccinating earlier in the 27 through 36-week period will maximize passive antibody transfer to the infant.
- The footnote for HPV vaccine has been updated to include the new 2-dose schedule for persons initiating the HPV vaccination series before age 15 years. In addition, bivalent

HPV vaccine has been removed from the schedule. This vaccine has been removed from the U.S. market, and all available vaccine doses have expired.

### Acknowledgments

Members of the Advisory Committee on Immunization Practices (ACIP) (current and past member rosters are available at (<https://www.cdc.gov/vaccines/acip/committee/members-archive.html>)); ACIP Child/Adolescent Immunization Work Group members: William L. Atkinson, MD, Harrisonville, Missouri; Katherine Brewer, MSN, Silver Spring, Maryland; Allison Kempe, MD, Aurora, Colorado; Susan Lett, MD, Boston, Massachusetts; Robin Liu, MD, Portland, Oregon; H. Cody Meissner, MD, Boston, Massachusetts; Amy B. Middleman, Oklahoma City, Oklahoma; Cynthia Pellegrini, Washington, DC; Diane Peterson, Saint Paul, Minnesota; José Romero, MD, Little Rock, Arkansas (Chair); Tina Simpson, MD, Birmingham, Alabama; Don Solimini, PA-C, Fishersville, Virginia; Rosemary Spence, MA, Denver, Colorado; Patricia Stinchfield, MPH, Saint Paul, Minnesota; Jennie Yoost, MD, Huntington, West Virginia. ACIP Child/Adolescent Immunization Work Group Contributors: Jennifer Hamborsky, MPH, Atlanta, Georgia; Lauren Hughes, MPH, Atlanta, Georgia; Suzanne Johnson-DeLeon, MPH, Atlanta, Georgia; David Kim, MD, Atlanta, Georgia; Andrew Kroger, MD, Atlanta, Georgia; Candice Robinson, MD, Atlanta, Georgia (CDC Lead); Raymond Strikas, MD, Atlanta, Georgia; Donna Weaver, MN, Atlanta, Georgia; Akiko Wilson, Atlanta, Georgia; Charles Wolfe, Atlanta, Georgia; JoEllen Wolicki, Atlanta, Georgia.

<sup>1</sup>Immunization Services Division, National Center for Immunization and Respiratory Diseases, CDC; <sup>2</sup>University of Arkansas for Medical Sciences and Arkansas Children’s Hospital, Little Rock, Arkansas; <sup>3</sup>Department of Pediatrics, University of Colorado Anschutz Medical Campus, Denver, Colorado; <sup>4</sup>March of Dimes, Washington, DC.

Corresponding author: Candice L. Robinson, [crobinson4@cdc.gov](mailto:crobinson4@cdc.gov), 404-718-1400.

### References

1. Grohskopf LA, Sokolow LZ, Broder KR, et al. Prevention and control of seasonal influenza with vaccines. MMWR Rep Recomm Rep 2016;65(No. RR-5).
2. Meites E, Kempe A, Markowitz LE. Use of 2-dose schedule for human papillomavirus vaccination—updated recommendations of the Advisory Committee on Immunization Practices. MMWR Morb Mortal Wkly Rep 2016;65:1405–8. <http://dx.doi.org/10.15585/mmwr.mm6549a5>
3. Updated 2016 ACIP statement on October 2016 hepatitis B vaccination recommendations (publication pending).
4. Briere EC. Food and Drug Administration approval for use of Hiberix as a 3-dose primary *Haemophilus influenzae* type b (Hib) vaccination series. MMWR Morb Mortal Wkly Rep 2016;65:418–9. <http://dx.doi.org/10.15585/mmwr.mm6516a3>
5. MacNeil JR, Rubin LG, Patton M, Ortega-Sanchez IR, Martin SW. Recommendations for use of meningococcal conjugate vaccines in HIV-infected persons—Advisory Committee on Immunization Practices, 2016. MMWR Morb Mortal Wkly Rep 2016;65:1189–94. <http://dx.doi.org/10.15585/mmwr.mm6543a3>
6. Updated ACIP statement on October 2016 meningococcal vaccination recommendations (publication pending).
7. Updated ACIP statement on October 2016 diphtheria and tetanus toxoids and acellular pertussis vaccination recommendations (publication pending).

## Advisory Committee on Immunization Practices Recommended Immunization Schedule for Adults Aged 19 Years or Older — United States, 2017

David K. Kim, MD<sup>1</sup>; Laura E. Riley, MD<sup>2</sup>; Kathleen H. Harriman, PhD<sup>3</sup>; Paul Hunter, MD<sup>4</sup>; Carolyn B. Bridges, MD<sup>1</sup>

On February 7, 2017, this report was posted as an MMWR Early Release on the MMWR website (<https://www.cdc.gov/mmwr>).

In October 2016, the Advisory Committee on Immunization Practices (ACIP) voted to approve the Recommended Adult Immunization Schedule for Adults Aged 19 Years or Older—United States, 2017. The 2017 adult immunization schedule summarizes ACIP recommendations in two figures, footnotes for the figures, and a table of contraindications and precautions for vaccines recommended for adults. These documents are available at <https://www.cdc.gov/vaccines/schedules>. The full ACIP recommendations for each vaccine can be found at <https://www.cdc.gov/vaccines/hcp/acip-recs/index.html>. The 2017 adult immunization schedule was also reviewed and approved by the American College of Physicians (<https://www.acponline.org>), the American Academy of Family Physicians (<https://www.aafp.org>), the American College of Obstetricians and Gynecologists (<http://www.acog.org>), and the American College of Nurse-Midwives (<http://www.midwife.org>).

A cover page has been added to the 2017 adult immunization schedule that contains information on select general principles pertinent to the adult immunization schedule, additional CDC resources, instructions for reporting vaccine adverse events related to vaccination and suspected cases of reportable

vaccine-preventable diseases, and an ACIP-approved list of standardized acronyms for vaccines recommended for adults.\* In addition, the table of contraindications and precautions for vaccines routinely recommended for adults, which was formerly a stand-alone document, has been incorporated into the adult immunization schedule.

### Changes in the 2017 Adult Immunization Schedule

Changes in the 2017 adult immunization schedule from the previous year's schedule include new or revised ACIP recommendations for influenza, human papillomavirus, hepatitis B, and meningococcal vaccines:

**Influenza vaccination (I).** Changes are related to the low effectiveness of the live attenuated influenza vaccine (LAIV) (FluMist, MedImmune) against influenza A(H1N1)pdm09 in the United States during the 2013–2014 and 2015–2016 influenza seasons and revised recommendations for the use of influenza vaccine among patients with egg allergy. These changes are reflected in the 2017 adult immunization schedule as follows:

- LAIV should not be used during the 2016–2017 influenza season.
- Adults with a history of egg allergy who have only hives after exposure to egg should receive age-appropriate inactivated influenza vaccine (IIV) or recombinant influenza vaccine (RIV).
- Adults with a history of egg allergy with symptoms other than hives (e.g., angioedema, respiratory distress, lightheadedness, or recurrent emesis, or who required epinephrine or another emergency medical intervention) may receive age-appropriate IIV or RIV. The selected vaccine should be administered in an inpatient or outpatient medical setting and supervised by a health care provider who is able to recognize and manage severe allergic conditions.

\* CDC encourages organizations that previously have relied on copying the adult immunization schedule on their websites to use syndication instead, as a more reliable method for displaying the most current and accurate adult immunization schedule. Use of content syndication requires a one-time step that ensures an organization's website displays the adult immunization schedule as soon as it is published or revised. The syndication code for the adult immunization schedule and instructions for its use can be found at <https://www.cdc.gov/vaccines/schedules/syndicate.html>. Requests for technical assistance for adult immunization schedule syndication can be sent to [ncirdwebteam@cdc.gov](mailto:ncirdwebteam@cdc.gov).

*Recommendations for routine use of vaccines in children, adolescents and adults are developed by the Advisory Committee on Immunization Practices (ACIP). ACIP is chartered as a federal advisory committee to provide expert external advice and guidance to the Director of the Centers for Disease Control and Prevention (CDC) on use of vaccines and related agents for the control of vaccine-preventable diseases in the civilian population of the United States. Recommendations for routine use of vaccines in children and adolescents are harmonized to the greatest extent possible with recommendations made by the American Academy of Pediatrics (AAP), the American Academy of Family Physicians (AAFP), and the American College of Obstetricians and Gynecologists (ACOG). Recommendations for routine use of vaccines in adults are harmonized with recommendations of AAFP, ACOG, and the American College of Physicians (ACP). ACIP recommendations approved by the CDC Director become agency guidelines on the date published in the Morbidity and Mortality Weekly Report (MMWR). Additional information about ACIP is available at <https://www.cdc.gov/vaccines/acip>.*

**Human papillomavirus vaccination (2).** Healthy adolescents who start their human papillomavirus (HPV) vaccination series before age 15 years are recommended to receive 2 doses of HPV vaccine. Adults and adolescents who did not start their HPV vaccination series before age 15 years should receive 3 doses of HPV vaccine. Changes in recommendations in the adult immunization schedule include updates regarding HPV vaccination for adults who did not complete the HPV vaccination series as adolescents. These changes are described in the 2017 adult immunization schedule as follows:

- Adult females through age 26 years and adult males through age 21 years who have not received any HPV vaccine should receive a 3-dose series of HPV vaccine at 0, 1–2, and 6 months. Males aged 22 through 26 years may be vaccinated with a 3-dose series of HPV vaccine at 0, 1–2, and 6 months.
- Adult females through age 26 years and adult males through age 21 years (and males aged 22 through 26 years who may receive HPV vaccine) who initiated the HPV vaccination series before age 15 years and received 2 doses at least 5 months apart are considered adequately vaccinated and do not need an additional dose of HPV vaccine.
- Adult females through age 26 years and adult males through age 21 years (and males aged 22 through 26 years who may receive HPV vaccine) who initiated the HPV vaccination series before age 15 years and received only 1 dose, or 2 doses less than 5 months apart, are not considered adequately vaccinated and should receive 1 additional dose of HPV vaccine.

**Hepatitis B vaccination (3).** The ACIP updated chronic liver disease conditions for which a hepatitis B vaccine (HepB) series is recommended. This change is described in the 2017 adult immunization schedule as follows:

- Adults with chronic liver disease, including, but not limited to, hepatitis C virus infection, cirrhosis, fatty liver disease, alcoholic liver disease, autoimmune hepatitis, and an alanine aminotransferase (ALT) or aspartate aminotransferase (AST) level greater than twice the upper limit of normal should receive a HepB series.

**Meningococcal vaccination (4,5).** There are two changes in meningococcal vaccination recommendations for 2017. First, the ACIP recommended that adults with human immunodeficiency virus (HIV) infection receive a 2-dose primary series of serogroups A, C, W, and Y meningococcal conjugate vaccine (MenACWY). Second, the ACIP provided updated dosing guidance for one of the serogroup B meningococcal vaccines (MenB) (MenB-FHbp [Trumenba, Pfizer]). Three doses of

MenB-FHbp should be administered at 0, 1–2, and 6 months to adults who are at increased risk for meningococcal disease, and those who are vaccinated during serogroup B meningococcal disease outbreaks. When MenB-FHbp is given to healthy adolescents and young adults who are not at increased risk for meningococcal disease, 2 doses of MenB-FHbp should be administered at 0 and 6 months (MenB-FHbp was previously recommended as a 3-dose series at 0, 2, and 6 months, consistent with the original vaccine licensure for this population). The dosing frequency and interval for the other MenB, MenB-4C (Bexsero, GlaxoSmithKline), have not changed: MenB-4C remains a 2-dose series, with doses administered at least 1 month apart. Either MenB vaccine can be used when vaccination is indicated. The change in ACIP recommendations on the use of MenB-FHbp does not imply a preference for one MenB over the other. These updates in meningococcal vaccination are reflected in the 2017 adult immunization schedule as follows:

- Adults with anatomical or functional asplenia or persistent complement component deficiencies should receive a 2-dose primary series of MenACWY, with doses administered at least 2 months apart, and be revaccinated every 5 years. They should also receive a series of MenB with either MenB-4C (2 doses administered at least 1 month apart) or MenB-FHbp (3 doses administered at 0, 1–2, and 6 months).
- Adults with HIV infection who have not been previously vaccinated should receive a 2-dose primary MenACWY vaccination series, with doses administered at least 2 months apart, and be revaccinated every 5 years. Those who previously received 1 dose of MenACWY should receive a second dose at least 2 months after the first dose. MenB is not routinely recommended for adults with HIV infection, because meningococcal disease in this population is caused primarily by serogroups C, W, and Y.
- Microbiologists who are routinely exposed to isolates of *Neisseria meningitidis* should receive 1 dose of MenACWY and be revaccinated every 5 years if the risk for infection remains, as well as either MenB-4C (2 doses administered at least 1 month apart) or MenB-FHbp (3 doses administered at 0, 1–2, and 6 months).
- Adults at risk because of a meningococcal disease outbreak should receive 1 dose of MenACWY if the outbreak is attributable to serogroup A, C, W, or Y; or, if the outbreak is attributable to serogroup B, either MenB-4C (2 doses administered at least 1 month apart) or MenB-FHbp (3 doses administered at 0, 1–2, and 6 months).

- Young adults aged 16 through 23 years (preferred age range is 16 through 18 years) who are healthy and not at increased risk for serogroup B meningococcal disease may receive either MenB-4C (2 doses administered at least 1 month apart) or MenB-FHbp (3 doses administered at 0, 1–2, and 6 months) for short-term protection against most strains of serogroup B meningococcal disease.

**Notable changes to Figures 1 and 2.** Changes in “Figure 1. Recommended immunization schedule for adults aged 19 years or older, by age group” and “Figure 2. Recommended immunization schedule for adults aged 19 years or older by medical condition and other indications” are as follows:

- In Figures 1 and 2, standardized acronyms for vaccines are used to promote simplicity and consistency, and their listing has been reordered. Ancillary information previously contained in the figures has been consolidated and moved to the cover page. Colored blocks instead of colored bars are used to denote indications. These figures must be used in conjunction with the footnotes, which contain important information for each vaccine and considerations for special populations.
- In Figure 2, the columns for medical conditions and other indications have been reordered to keep medical conditions together and special populations together. Additional footnotes mark appropriate columns of medical conditions and other indications to refer the reader to view relevant vaccine-specific information.
- In Figure 2, the color of the indication block for MenACWY for HIV infection has been changed to yellow (recommended for adults who meet the age requirement, lack documentation of vaccination, or lack evidence of past infection) from purple (recommended for adults with additional medical conditions or other indications).

#### Changes to footnotes.

- Footnotes are limited to the information pertaining to vaccines listed in Figures 1 and 2 and are organized by vaccine-specific information and considerations for special populations (e.g., pregnant women and adults with HIV infection). The footnote labeled “additional information,” contained in previous versions of the adult immunization schedule, has been moved to the cover page. The footnote related to immunocompromising conditions has been removed, but vaccine-specific information on immunocompromising conditions has been added to the appropriate footnotes (e.g., the footnote for pneumococcal vaccination).
- The format for the footnotes has been condensed, simplified, and standardized. The format for pneumococcal;

human papillomavirus; meningococcal; varicella; and measles, mumps, and rubella vaccination footnotes has undergone substantial revision.

**Other changes.** Lastly, the table of contraindications and precautions for vaccines routinely recommended for adults, which previously was a stand-alone document, has been incorporated into the adult immunization schedule. The content of the table has been consolidated and simplified.

#### More Information

Details on these updates and information on other vaccines recommended for adults are available online under Adult Immunization Schedule, United States, 2017 (<https://www.cdc.gov/vaccines/schedules/hcp/adult.html>) and in the *Annals of Internal Medicine* (6). The full ACIP recommendations for each vaccine are also available online (<https://www.cdc.gov/vaccines/hcp/acip-recs/index.html>).

#### Acknowledgments

Advisory Committee on Immunization Practices (ACIP member rosters are available online at <https://www.cdc.gov/vaccines/acip/committee/members-archive.html>); ACIP Adult Immunization Work Group.

<sup>1</sup>Immunization Services Division, National Center for Immunization and Respiratory Diseases, CDC; <sup>2</sup>Harvard University; <sup>3</sup>California Department of Public Health; <sup>4</sup>University of Wisconsin.

Corresponding author: David K. Kim, [dkim@cdc.gov](mailto:dkim@cdc.gov), 404–639–0969.

#### References

1. Grohskopf LA, Sokolow LZ, Broder KR, et al. Prevention and control of seasonal influenza with vaccines. *MMWR Recomm Rep* 2016;65(No. RR-5). <http://dx.doi.org/10.15585/mmwr.rr6505a1>
2. Meites E, Kempe A, Markowitz LE. Use of 2-dose schedule for human papillomavirus vaccination—updated recommendations of the Advisory Committee on Immunization Practices. *MMWR Morb Mortal Wkly Rep* 2016;65:1405–8. <http://dx.doi.org/10.15585/mmwr.mm6549a5>
3. Updated 2016 ACIP statement on October 2016 hepatitis B vaccination recommendations (publication pending).
4. MacNeil JR, Rubin LG, Patton M, Ortega-Sanchez IR, Martin SW. Recommendations for use of meningococcal conjugate vaccines in HIV-infected persons—Advisory Committee on Immunization Practices, 2016. *MMWR Morb Mortal Wkly Rep* 2016;65:1189–94. <http://dx.doi.org/10.15585/mmwr.mm6543a3>
5. Updated ACIP statement on October 2016 meningococcal vaccination recommendations (publication pending).
6. Kim DK, Riley LE, Harriman KH, Hunter P, Bridges CB. Advisory Committee on Immunization Practices. Recommended immunization schedule for adults aged 19 years or older, United States, 2017. *Ann Intern Med* 2017;166:209–18. <http://annals.org/aim/article/doi/10.7326/M16-2936>